

Sensory Symptoms Checklist

The following checklist is not a diagnostic tool. Rather, it is an indicator of sensory overor under-responsiveness. The purpose of this tool is to assist in developing an appropriate treatment plan, and/or sensory diet, for an individual with sensory modulation difficulties.

Remember that it is normal for all people to be sensitive to certain sensations. Sensory modulation becomes a disorder only when it is negatively impacting a person's life (i.e. ability to pay attention, learn, socialize, relax).

* Indicates Sensory-Seeking behavior. Sensory-Seeking is a form of underresponsiveness but these individuals seek intense sensation to make up for the underresponsiveness of their nervous systems. The general under-responders tend to be more passive and sedentary than the sensory-seekers, but still share many symptoms.

TACTILE (TOUCH)

Symptoms of Over-Responsiveness:	Symptoms of Under-Responsiveness:
avoids affectionate touch	always touching others*
bothered by seams in clothing,	as a young child, may prefer to
tags, waistbands, etc.	be without clothes and barefoot
avoids messy play including	doesn't seem to notice messy
finger-painting and play-doh	hands or face
dislikes nail-trimming/hair-cutting	touches everything, brings objects
hair-brushing/tooth-brushing	and toys to mouth frequently*
limited food preferences,	may stuff too much food in mouth
sensitive to food textures	seems unaware of light touch
excessively ticklish	doesn't seem to notice cuts and
avoids standing close to others	scrapes, doesn't mind shots
startles easily when touched	drools, doesn't wipe runny nose
unexpectedly	unintentionally rough on pets or
overreacts to minor injuries or	other kids
mosquito bites	craves intense flavors – salty, sweet sour, spicy*
	difficulty with fine motor tasks



PROPRIOCEPTIVE (PRESSURE ON MUSCLES AND JOINTS)

Since proprioceptive input is always helpful to the nervous system, we do not see <i>over-responsivity</i> in this category.	Symptoms of Under-Responsiveness: loves jumping, climbing, wrestling, and crashing activities* walks on toes*, or walks heavily difficulty with fine motor tasks prefers crunchy and chewy foods* sucks thumb or fingers, chews on clothes and toys grinds teeth, cracks knuckles* loves tight hugs and "squishing" activities or positions* chooses thick or heavy blankets self-abusive behaviors — pinching, biting, head-banging* prefers tight-fitting clothes or may wear a snug jacket constantly* uses too much or too little force on objects (juice boxes, crayons)
VESTIBULAR (MOVEMENT) Symptoms of Over-Responsiveness: as a baby, dislikes being held away from adult's body or tossed in air, may be "clingy" avoids swings, slides, anything that requires feet to be off the ground fearful of escalators, elevators, and all heights gets motion sick easily dislikes leaning back for hair- washing or floating on back in pool	Symptoms of Under-Responsiveness: as a baby, needs to be rocked a lot, is happiest in stroller, swing, bouncer* dislikes sedentary tasks* is in constant motion, loves spinning, swinging, being upside down* may have low muscle tone (muscles and joints seem too soft and floppy) "W" sitting on floor, slumps, leans in chair loves spinning, loves amusement park rides, is a "thrill-seeker"* rocks self or moves head back and forth while sitting*



INTEROCEPTION (INTERNAL SENSATIONS)	
Symptoms of Over-Responsiveness:	Symptoms of Under-Responsiveness:
seems to have a low pain	difficult to toilet-train
tolerance	apparent lack of hunger/thirst
displays significant distress	high pain tolerance or shows
when cold after bath or after	inappropriate response to pain
stepping outside on hot day	difficulty falling asleep/staying asleep
extreme fear of medical procedures	doesn't dress appropriately for
frequent complaints of various	weather
aches and pains	doesn't seem aware of symptoms of
	illness such as sore throat or nausea
AUDITORY (SOUND)	
Symptoms of Over-Responsiveness:	Symptoms of Under-Responsiveness:
as a baby, startled easily, cries with	may not consistently respond to
vacuum, hair-dryers, toilet flushing	name (rule out hearing loss)
dislikes noisy places	prefers t.v. and music to be loud*
easily distracted and bothered	often doesn't notice background
by background noises	noises
cries, covers ears with loud or	makes own sounds frequently,
unexpected sounds	enjoys silly sounds of others*
asks others to be quiet	says "What?" frequently even though
	hearing is intact
	may appear oblivious to some
	sounds, has difficulty locating sound
	may use self-talk to get through a
	task
VISUAL Symptoms of Over-Responsiveness:	Symptoms of Under-Responsiveness:
bothered by bright lights, sun	loves shiny, spinning or moving
dislikes visually busy places	objects*
(stores, playgrounds, cluttered	difficulty with eye-hand coordination
rooms)	tasks (catching a ball, stringing
avoids eye contact	(beads, tracing and writing)
prefers dim lighting, shade	loves action-packed, colorful
rubs eyes a lot, may get	t.v. shows and electronic games*
headaches from reading	difficulty distinguishing between
	similar letters and shapes
	difficulty with visual-tracking, may
	lose place frequently while reading



GUSTATORY AND OLFACTORY (TASTE AND SMELL)

Symptoms of Over-Responsiveness:	Symptoms of Under-Responsiveness:
talks about smell a lot, notices odors others don't notice	smells and licks inedible objects such as toys and play-doh*
plugs nose, avoids places with strong smells (some restaurants) dislikes new foods, may have very limited food preferences	likes strong odors such as perfume cleaning products, and gasoline* prefers strongly flavored foods - spicy, salty, bitter, sour, sweet*

^{*}Indicates sensory-seeking behavior. See note at top of form.